



Bayer CropScience

Bayer CropScience LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

Folicur® 3.6 F Foliar Fungicide

EPA Reg. No. 264-752

For Use of Specified Diseases on Various Crops

Supplemental Label

Folicur® 3.6 F Foliar Fungicide

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read this label and the product package label before using this product. This Supplemental Label must be in possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for Folicur® 3.6 F Foliar Fungicide.

USE DIRECTIONS FOR SPECIFIC CROPS

RECOMMENDED APPLICATIONS

CROP	DISEASE	RATE OF FOLICUR 3.6 F
Asparagus	Rust (Puccinia spp.)	4 to 6 fl. oz. per acre
	<p>Notes: Apply FOLICUR 3.6 F as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl oz of FOLICUR 3.6 F per acre (0.11 lb ai – 0.17 lb ai per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14-day interval as necessary to maintain control of rust. Do not apply to harvestable spears. Do not apply within 100 days of harvest in California and 180 days in all other states. Do not make more than three foliar applications per season (18 fl oz/acre or 0.51 lb ai/acre).</p>	
<p>General Comments: Applications may be made using ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with FOLICUR 3.6 F. FOLICUR 3.6 F is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating FOLICUR 3.6 F with other DMI fungicides may lead to resistance.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF FOLICUR 3.6 F
Beans (fresh & dry except succulent shelled)	Rust (<i>Uromyces appendiculatus</i>)	4 to 6 fl. oz. per acre
	Notes: Apply FOLICUR 3.6 F in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control. Beans, fresh: FOLICUR 3.6 F may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of FOLICUR 3.6 F per acre per crop season. Beans, dry: FOLICUR 3.6 F may be applied up to 14 days before harvest. Do not apply more than 12 fl. oz. of FOLICUR 3.6 F per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with FOLICUR 3.6 F. FOLICUR 3.6 F must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, FOLICUR 3.6 F will be resistant to weathering. FOLICUR 3.6 F is a demethylation inhibitor (DMI) fungicide (Group 3).		
Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF FOLICUR 3.6 F
Corn (sweet corn, field corn, field corn grown for seed, and popcorn)	Rust (<i>Puccinia spp.</i>)	4 to 6 fl. oz. per acre
	Northern leaf blight (<i>Helminthosporium turcicum</i>)	
Southern leaf blight (<i>Helminthosporium maydis</i>)		
Northern leaf spot (<i>Helminthosporium carbonum</i>)		
Gray leaf spot (<i>Cercospora zea-maydis</i>)		
Notes: Apply FOLICUR 3.6 F in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. A maximum of 24 fl. oz. (1.5 pint) of FOLICUR 3.6 F may be applied per acre per crop season. Sweet corn: FOLICUR 3.6 F may be applied up to 7 days before the harvest of ears or forage, and 49 days before the harvest of fodder. Field, seed or popcorn: FOLICUR 3.6 F may be applied up to 21 days before the harvest of forage, and 36 days before the harvest of grain or fodder.		
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with FOLICUR 3.6 F. FOLICUR 3.6 F must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, FOLICUR 3.6 F will be resistant to weathering. FOLICUR 3.6 F is a demethylation inhibitor (DMI) fungicide (Group 3).		
Restricted-entry interval (REI) for sweet corn = 19 days.		
Restricted-entry interval (REI) for all corn except sweet corn = 12 hours.		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF FOLICUR 3.6 F
Cotton	Southwestern cotton rust (<i>Puccinia cacabata</i>)	6 to 8 fl. oz. per acre
	Notes: Apply FOLICUR 3.6 F in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. FOLICUR 3.6 F may be applied up to 30 days before harvest. Do not apply more than 24 fl. oz. of FOLICUR 3.6 F per acre per crop season.	
<p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with FOLICUR 3.6 F. FOLICUR 3.6 F must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, FOLICUR 3.6 F will be resistant to weathering. FOLICUR 3.6 F is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF FOLICUR 3.6 F
Cucurbit Vegetables Group Chayote Chinese waxgourd Citron melon Cucumber Gherkin Edible gourd, (includes hyotan, cucuzza, hechima and Chinese okra) Momordica spp. (includes balsam apple, balsam pear, bitter melon and Chinese cucumber) Muskmelon (includes cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon) Pumpkin Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini) Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash) Watermelon	Powdery mildew (<i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i>) (<i>Erysiphe cichoracearum</i>)	4 to 6 fl. oz. per acre
	Gummy stem blight - suppression (<i>Didymella bryonae</i>) (watermelon, squash, pumpkin, and melons only)	8 fl. oz. per acre
<p>Notes: Apply the specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals. FOLICUR 3.6 F may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of FOLICUR 3.6 F per acre per crop season.</p>		
<p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with FOLICUR 3.6 F. FOLICUR 3.6 F must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, FOLICUR 3.6 F will be resistant to weathering. FOLICUR 3.6 F is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF FOLICUR 3.6 F
Dry bulb onion Garlic Great-headed (elephant) garlic Welch onion Shallot	White rot (Sclerotium cepivorum)	White rot: 20.5 fl oz per acre applied in a 4 to 6 inch band over/into each furrow. May be applied by chemigation to control white rot.
	Rust (Puccinia allii, Puccinia porri) Purple blotch (Alternaria porii)	4 to 6 fl. oz per acre
	<p>White rot: For the control of white rot, make one application in the furrow at the time of planting. The in-furrow application should be made at the rate of 20.5 fl. oz Folicur 3.6 F per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 to 6 fl oz/acre.</p> <p>Rust: For the control of rust make foliar applications at the rate of 4 to 6 fl. oz Folicur 3.6 F per acre per application. Repeat at an interval of 10 to 14 days. Apply FOLICUR 3.6 F in a protective spray schedule or when weather conditions are favorable for rust development.</p> <p>Notes: Do not apply more than 32.5 fl. oz. Folicur 3.6 F per acre per season if an in-furrow treatment is made. If Folicur 3.6 F is not applied as an in-furrow treatment then do not apply more than 12 fl oz. Folicur 3.6 F per acre per season as a foliar spray. Do not apply within 7 days of harvest (PHI = 7 days).</p>	
<p>General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with FOLICUR 3.6 F. FOLICUR 3.6 F must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, FOLICUR 3.6 F will be resistant to weathering. FOLICUR 3.6 F is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF FOLICUR 3.6 F
Green onion Leek Spring onion Scallion Japanese bunching onion Green shallots Green eschalots	White rot (<i>Sclerotium cepivorum</i>) suppression only Rust (<i>Puccinia allii</i> , <i>Puccinia porri</i>) Purple blotch (<i>Alternaria porii</i>)	4 to 6 fl. oz per acre
<p>For the control of diseases make foliar applications using an interval of 10 to 14 days. Apply FOLICUR 3.6 F in a protective spray schedule or when weather conditions are favorable for rust development.</p> <p>Notes: Do not apply more than 24 fl. oz. Folicur 3.6 F per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).</p>		
<p>General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with FOLICUR 3.6 F. FOLICUR 3.6 F must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, FOLICUR 3.6 F will be resistant to weathering. FOLICUR 3.6 F is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF FOLICUR 3.6 F
Hops	Powdery mildew (<i>Sphaerotheca humuli</i> / <i>Sphaerotheca macularis</i>)	4 to 8 fl. oz. per acre
<p>Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. FOLICUR 3.6 F may be applied up to 14 days before harvest. Do not apply more than 32 fl. oz. of FOLICUR 3.6 F per acre per crop season. Increase the spray volume and the application rate as vine growth increases during the season.</p>		
<p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with FOLICUR 3.6 F. FOLICUR 3.6 F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, FOLICUR 3.6 F will be resistant to weathering. FOLICUR 3.6 F is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF FOLICUR 3.6 F
Leafy Brassica Greens Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuma Mustard greens Mustard spinach Rape greens Turnip greens	Cercospora leaf spot (Cercospora brassicicola)	3 to 4 fl. oz per acre
	Powdery mildew (Erysiphe cruciferarum)	
	Alternaria leaf spot (Alternaria brassicicola)	
Notes: Do not apply more than 16 fl. oz. Folicur 3.6 F per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).		
General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with FOLICUR 3.6 F. FOLICUR 3.6 F must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, FOLICUR 3.6 F will be resistant to weathering. FOLICUR 3.6 F is a demethylation inhibitor (DMI) fungicide (Group 3). Restriction: Application to turnip greens is limited to East of the Rockies Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF FOLICUR 3.6 F
Garden beet roots and tops (leaves)	Cercospora leaf spot (Cercospora beticola)	3 to 7.2 fl. oz per acre
	Notes: Make applications on a 14 day intervals. Do not apply more than 28.8 fl. oz. Folicur 3.6 F per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).	
General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank-mixed with FOLICUR 3.6 F. FOLICUR 3.6 F must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, FOLICUR 3.6 F will be resistant to weathering. FOLICUR 3.6 F is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF FOLICUR 3.6 F
Lychee	Anthracnose (<i>Colletotrichum gloeosporioides</i>)	4 to 6 fl. oz. per acre
	Notes: Begin first application of FOLICUR 3.6 F as panicle emerges. Spray up to 6 fl. oz. per acre every 10 days thereafter for a total of 8 sprays. Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only. Do not apply more than 48 fl. oz. of FOLICUR 3.6 F per acre per season. FOLICUR 3.6 F can be applied up to and including the day of harvest (PHI = 0 days).	
<p>General Comments: For optimum disease control, the lowest labeled rate of a non-ionic spray surfactant should be tank-mixed with FOLICUR 3.6 F. FOLICUR 3.6 F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, FOLICUR 3.6 F will be resistant to weathering. FOLICUR 3.6 F is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 2 days.</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF FOLICUR 3.6 F
Mango (Post-Harvest)	Anthracnose (<i>Colletotrichum gloeosporioides</i>)	7.5 fl. oz. per 100 gal. water / 50,000 lbs. fruit
	Notes: FOLICUR 3.6 F can be applied as a post-harvest treatment by mixing at the specified rate. Application to the fruit is made by spraying the dilute FOLICUR 3.6 F solution onto brushes in a commercial packing line.	

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF FOLICUR 3.6 F
Okra	Cercospora leaf spot (<i>Cercospora spp.</i>)	4 to 6 fl. oz. per acre
	Notes: Apply specific dosage of FOLICUR 3.6 F in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Applications may be made no closer than 3 days before harvest. Do not apply more than 24 fl. oz. of FOLICUR 3.6 F per acre per season.	
<p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with FOLICUR 3.6 F. FOLICUR 3.6 F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, FOLICUR 3.6 F will be resistant to weathering. FOLICUR 3.6 F is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF FOLICUR 3.6 F
Pecan	Brown leaf spot <i>(Sirosporium diffusium)</i> Downy spot <i>(Mycosphaerella caryigena)</i> Liver spot <i>(Gnomonia caryae)</i> Scab <i>(Cladosporium caryigenum)</i> Vein spot <i>(Gnomonia nerviseda)</i> Zonate leaf spot <i>(Grovesinia pyramidalis)</i>	4 to 8 fl. oz. per acre
	<p>Notes: Apply FOLICUR 3.6 F in a preventive spray schedule beginning at early bud break (young leaves unfolding), and continue applications at 10- to 14-day intervals through the pollination period. FOLICUR 3.6 F should be applied at 4 fl. oz. per acre in a tank-mix with the recommended rate of Super-Tin® in cover sprays. Follow label directions for the use of Super-Tin. Do not add a surfactant to the spray solution when tank-mixing FOLICUR 3.6 F with Super-Tin. Apply FOLICUR 3.6 F in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 to 8 fl. oz. per acre of FOLICUR 3.6 F to full-size mature trees, and 4 to 6 fl. oz. per acre of FOLICUR 3.6 F to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases, or when severe disease conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases. Do not apply after shucks begin to split. A maximum of 32 fl. oz. of FOLICUR 3.6 F may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.</p>	
<p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with FOLICUR 3.6 F. FOLICUR 3.6 F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, FOLICUR 3.6 F will be resistant to weathering. FOLICUR 3.6 F is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE(S)	RATE OF FOLICUR® 3.6F
Soybean	Rust (<i>Phakopsora pachyrhizi</i>) Powdery mildew (<i>Microsphaera diffusa</i>)	3 to 4 fl oz per acre
<p>Use Directions: Apply Folicur® 3.6 F as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use of the higher rates and shorter spray intervals are recommended when disease pressure is severe. The lowest label recommended rate of a spray surfactant must be tank-mixed with Folicur® 3.6F. Folicur® 3.6F should be applied in a minimum for 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.</p> <p>Restrictions: Applications may not be made within 21 days of harvest. Do not apply more than 3 applications per season. Do not apply more than 12 fl. oz/a per use season.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF FOLICUR 3.6 F
Sunflower	Rust (<i>Puccinia helianthi</i>)	4 to 6 fl. oz. per acre
	<p>Notes: Apply specific dosage of FOLICUR 3.6 F at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Application may be repeated at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Do not apply more than 16 fl. oz. of FOLICUR 3.6 F per acre per season or within 50 days of harvest.</p>	
<p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with FOLICUR 3.6 F. Contact your state Extension Service or Bayer representative for a list of approved surfactants. FOLICUR 3.6 F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, FOLICUR 3.6 F will be resistant to weathering. FOLICUR 3.6 F is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF FOLICUR 3.6 F
Turnip (Application is limited to East of the Rockies)	Cercospora leaf spot (<i>Cercospora brassicola</i>)	4 to 7.2 fl. oz. per acre
	<p>Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals. FOLICUR 3.6 F may be applied up to 7 days before harvest. Do not apply more than 28.8 fl. oz. of FOLICUR 3.6 F per acre per crop season.</p>	
<p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with FOLICUR 3.6 F. FOLICUR 3.6 F must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, FOLICUR 3.6 F will be resistant to weathering. FOLICUR 3.6 F is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

Folicur is a registered trademark of Bayer.