Extension Report No. 9

1990 SUNFLOWER TROWER SURVEY

of Pest Problems and Pesticide Use in North Dakota

H. A. Lamey, D. K. McBride, R. K. Zollinger, J. L. Luecke, M. P. McMullen, and D. R. Berglund



NDSU EXTENSION SERVICE

FEBRUARY 1992

Procedures

This is the first survey of North Dakota sunflower grower pest problems and pesticide use. A single page survey form was designed with questions on pest problems, pesticide use, and alternative or non-chemical control practices (Figure 1). Every third name was selected from a grower mailing list of over 12,000 North Dakota sunflower growers. Survey forms were mailed on November 9, 1990 to 4,288 sunflower growers. A self-addressed stamped envelope was enclosed for returning the survey form. The survey asked respondents to identify: county where sunflowers grown; acres of oilseed and/or confection hybrids planted; major production problems encountered; major insect, disease and weed problems; percent bird damage; pesticides used; injury and control from the use of herbicides; and use of integrated pest management (IPM) and other alternative pest control techniques.

Identification of counties where production occurred was used to apportion the data among the nine reporting districts of the North Dakota Agricultural Statistics Service (1), as shown in Figure 2.

RESULTS

Approximately one fourth (1,015) of the mailed forms were returned. Of those returned, 363 had no data; the other 652, or 15%, had useable data. These 652 growers planted 189,204 acres, which is 14% of the total of 1,370,000 acres planted in North Dakota (1). Respondents planted 122,636 acres of oilseed hybrids, or 13% of the 980,000 acres planted statewide, and 66,568 acres of confection hybrids, or 17% of the 390,000 acres planted statewide.

Counties Leading in Sunflower Acreage. The eight leading counties among the respondents and the percent of respondents' acres planted in each were: Stutsman, 12%; Barnes, 9%; Wells, 6%; LaMoure, 6%; Cass, 6%; Foster, 5%; Ransom, 5%; and Benson, 5%. These eight counties accounted for 53% of the total acres planted by respondents (Table 1).

Production Problems. Insects were the worst production problem for 23% of respondents, diseases for 19%, weather for 17%, bird damage for 16%, and weeds for 12% (Table 2). No differences were reported




Figure 2. Crop reporting districts of the North Dakota Agricultural Statistics Service.

Mag Carcle one): 1990 (ARMK 1-3, 1 = MORST) 6 Prov. (spring) 1 2 3 1 2 3 1 2 3 3 2 3 3 3 3 3 3	otal acres planted in 1990		EVALUATE WEED CON	NTROL AND SUNF	LOWER INJURY		
Ounty where grown Acres county. Name and the second team one county where grown in more than one county, list each county and acres. Acres 3	Acres planted to oilgood		Mark weed cont	crol used an	di .		
CHIEBUL INJURY (If sunflower grown in more than one county, list each county and sorres) Acres (If sunflower grown in more than one county, list each county and sorres) Acres County and sorres) County and sorres County and	Acres planted to confect	nyorida on hubrida	indicate acres		r		
Culty Vales grown in acre than one county, list each county and acres) (If sunflower grown in acre than one county, list each county and acres) (If sunflower grown in acre than one county, list each county and acres) (If sunflower grown in acre than one county, list each county and acres) (If sunflower grown in acre than one county, list each county and acres) (If sunflower grown in acre than one county, list each county and acres) (If sunflower grown in acre than one county, list each county and acres) (If sunflower grown in acre than one county, list each county and acres) (If sunflower grown in acre than one county, list each county and acres) (If sunflower grown in acre than one county, list each county and acres) (If sunflower grown in acres) (If su	Marao bigured to courect	on hyprids	each item. Co				
(if sunflower grown in more than one county, list each county and acres) County and acres) County and acres) Acres Defair Production (Acres Defair Condition (Acres Defair County) and acres) REATEST PRODUCTION REATEST PROD	ounty where grown	ACTOS	application, o	louble			
County and acres 1-past			as double acre				
	(if sunflower grown in more	than one county, list each	de double acte	2	Acres		
1.	county and acres)		Weed control u	sed			
REATES PRODUCTION ORDICIT MINISTRUCER MISSO (Little one): 1190 (MARK 1-3, 1 - WORST) ORDICITAL SHOPPONE 123						1 2 3 4	1 2 3 4
### WASST DILEASE PROBLEMS IN 4 Sonalan	200 To 10	8,	2. Eptam/Gene	p (fall)	Name and the same		
Mone	REATEST PRODUCTION		1. Eptam/Gene	p (spring)			
March Marc	ROBLEM IN SUNFLOWER	WORST DISEASE PROBLEMS IN	5. Sonalan +	Entam/Genen			
None	N 1990 (circle one):	1990 (RANK 1-3, 1 - WORST)	6. Prowl (fal	11			
Scientinia will S. Treilan (fail) 12 14	one	None	7 Proul (enr	ingl			
Rust 10. Terlian Eptam/Genep 1 2 3 4 1		Sclerotinia wilt	8. Treflan (f	a11)			
Phona black Stem 11. Lesso		Sclerotinia head rot	9. Treflan (s	pring)			1 2 3 4
### SECTION Content of the content o		Rust	10. Treflan +	Eptam/Genep			
ind damage		Proma black stem				20 10 10 10 10 10 10 10 10 10 10 10 10 10	
WORST HEETED PROBLEMS IN 1930 (RANK 1-3, 1 = 1) = 1930 (RANK 1-3, 1 = 1) = 1930 (RANK 1-3, 1 = 1930 (RANK 1-3, 1 = 1) = 1					-		
WORST HEETED PROBLEMS IN 1930 (RANK 1-3, 1 = 1) = 1930 (RANK 1-3, 1 = 1) = 1930 (RANK 1-3, 1 = 1930 (RANK 1-3, 1 = 1) = 1	ther (specify)		14. Gramoxone	Extra	Control of the Contro	1 2 3 4	1 2 1 4
WORST HEETED PROBLEMS IN 1930 (RANK 1-3, 1 = 1) = 1930 (RANK 1-3, 1 = 1) = 1930 (RANK 1-3, 1 = 1930 (RANK 1-3, 1 = 1) = 1	- 10 To 10 T		15. Leafex-3/D	efol		1 2 3 4	1 2 3 4
WORST HEETED PROBLEMS IN 1930 (RANK 1-3, 1 = 1) = 1930 (RANK 1-3, 1 = 1) = 1930 (RANK 1-3, 1 = 1930 (RANK 1-3, 1 = 1) = 1			16. No herbici	de used	Name of the last o	1 2 3 4	1 2 3 4
Sed (RANK 1-3, 1 and 1 a						1 2 3 4	1 2 3 4
Mone 1234 12	ORST WEED PROBLEMS IN	WORST INSECT PROBLEMS IN					
None Seed weevil Banded sunflower moth Stem weevil Sunflower beatle Sunflower head moth Sunflower head moth Sunflower head moth Sunflower moth Sunflower head moth Sunflower moth Sunflower head moth Sunflower	990 (RANK 1-3, 1 -	1990 (RANK 1-3, 1 - WORST)	19. Others (sp	ecity)			
Seed vee'vil		None				1 2 1 4	
Stem weevil Stem weevil Stem weevil Fortail (Pigeon grass) Sunflower beatle Sunflower beatle Sunflower head moth Sunflower midge Sunflower		Seed weevil	magazine say a 4 a				
Foxtail (Pigeon grass) Wild buckwheat Wild mustard Sunflower midge Grasshopper Wild cats Other (specify) Other (specify) Other (specify) Other (specify) No. Acres No. of BIRD FROBLEMS AND LOSSES: Acres Klyl parathion 1		Banded sunflower moth					
Wild buckwheat Wild mustard Sunflower midge Redroot pigweed Wild oats Other (specify) Other (specify) Nother (specify)	Fourtail (Dianes	Stem weevil					
Redroot pigweed Grasshopper Other (specify) Other (specify) Other (specify) Other (specify) Other (specify) Other (specify) No. Acres No. of BIRD PROBLEMS AMD 10SSES: Acres Traited Spraya Mark which best fits 15-10/acre Striyl parathion (\$ yteld loss) 10 parathion 15-10t 100 parathion 15-10t 100 per other (specify) Indiane/Maneb (seed treatment) Indiane/Maneb (seed treatment) No. Acres No. of Funcicles USED ON SUMFLOMER IN 1990: Acres No. of Funcicles Treated Spraya Mark which best fits 15-10/acre 150-100 per other	Wild buckwheat	Sunflower beetle					
Redroot pigueed wild oats Other (specify) Other (specify) Other (specify) SECTICIDE(s) USED ON WINDER IN 1990. No. Acres No. of BIRD PROBLEMS AND LOSSES: Mark which best fits Style parathion Sans XL 10-15 Acres Solution Solu	Wild mustard	Sunflower midde					
Other (specify) Nother (specify) No. Section of the specify of the section of th	Redroot pigweed	Grashopper	*				
Other (specify) Secticide	Wild oats	Other (specify)		INTEGRATED PES	T MANAGEMENT		
SECTICIDE(s) USED ON INFLORER IN 1990: NO. INFLORER IN 1990: NO. INSECTICIDE(s) USED ON INFLORER IN 1990: NO. INFLORER IN 1990: INFLORER IN	Other (specify)	Name of the second seco					
If yes, how many acres? If yes, estimate economic return from using consultant: 1				scout sunfl	war?	es no	
INFLOWER IN 1990: No. Acres No. of BIRD PROBLEMS AND 1055E5: Secticide Treated Sprays Mark which best fits 15-10/acre thyl parathion 15-10% Sens NL 10-25% Sens NL 10-25% Servin XLR plus 12-5-50% Indiane/Maneb (seed treatment) Cher (specify) Results of survey will be published in The Sunflower thank you. Please return by November 25, 1990 Results of survey will be published in The Sunflower thank you. Please return by November 25, 1990 If yes, estimate economic return from using consultant: 12-20 + /acre				If yes, how	many acres	7	
No. No. Acres No. of BIRD PROBLEMS AND LOSSES: S10-15/acre S15-20/acre S10-15/acre				If yes, est	imate econo	mic	
No. Acres No. of BIRD PROBLEMS AND LOSSES: S10-15/acre S51-16/acre S11-16/acre		96				ultant:	
Acres No. of BIRD PROBLEMS AND 1055E5: specificide Treated Sprays thyl parathion				\$20 + /a	CIO		
Mark which best fits (* yield loss)				\$15-20/a	270		G.
thyl parathion (* yield loss)				S5=10/ac			*
So				50-5/acr			
		(* yield loss)		so	(2)		
Sana XL Dorsban 4E Dorsban 4D Dor	-3 parathion			other			
Josepha 4E	sana XL	10-25%		CONTRACTOR OF THE STATE OF THE			
pesticides: print XLR plus Gradan 15G Gradan 15G Gradan 15G Gradan 15G Gradan 15G Greet treatment) FUNGICIDES USED ON SUNFIGNER IN 1990: Cher (specify) No. Acres No. of Fungicide Treated Sprays Dithane Apron (seed treatment) None used (list acres) Other (specify) Results of survey will be published in The Sunflower Chank you. Please return by November 25, 1990 Acres No. of Fundicide Treated Sprays Other (specify) Acres No. of Fundicide Treated Sprays Other (specify) Acres Other (specify)	orsban 4E	25-50%			ther than		
Preceding crop How long since previous sunflower? FUNGICIDES USED ON SUNFLOWER IN 1990: No. Acres No. of Fungicide Treated Sprays Dithane Apron (seed treatment) None used (list acres) Other (specify) Results of survey will be published in The Sunflower Chank you. Please return by November 25, 1990 ACRES NO. OF Fungicide Treated Sprays Dithane Apron (seed treatment) None used (list acres) Other (specify) ACRES NO. OF Fungicide Treated Sprays Dithane Apron (seed treatment) None used (list acres) Other (specify) ACRES NO. OF Fungicide Treated Sprays Dithane Apron (seed treatment) None used (list acres) Other (specify) ACRES NO. OF Fungicide Treated Sprays Dithane Apron (seed treatment) None used (list acres) Other (specify) ACRES NO. OF Fungicide Treated Sprays Dithane Apron (seed treatment) None used (list acres) Other (specify) ACRES NO. OF Fungicide Treated Sprays Dithane Apron (seed treatment) None used (list acres) Other (specify) ACRES NO. OF TIMES NO	uradan 4F	50-100%		pesticides:			
indame/Maneb (seed treatment) one used (list acres) SUNFLOWER IN 1990: No. Acres No. of Fundicide Treated Sprays Dithane Apron (seed treatment) None used (list acres) Other (specify) Results of survey will be published in The Sunflower Chank you. Please return by November 25, 1990 Acres No. of Fundicide Treated Sprays Other (specify)		E CONTROL CONT		Crop rotation	onacre	S	
(seed treatment) one used (list acres) There (specify) FUNGICIDES USED ON SUNFLOWER IN 1990: No. Acres No. of Fungicide Treated Sprays Dithane Apron (seed treatment) None used (list acres) Other (specify) Results of survey will be published in The Sunflower Thank you. Please return by November 25, 1990 Author Ambur Amuly Arthur Lamey		452. 720		How long of	op	E 10	
No. Resistant Varietyacres				sunflower?	rea braaton		
No. Resistant Varietyacres	one used (list acres)			Cultivation	(No.	of times)	
Acres No. of Fungicide Treated Sprays Dithane Apron (seed	ther (specify)			Resistant Va	riety	acres	
Fungicide Treated Sprays Dithane Apron (seed treatment) None used (list acres) Other (specify) Results of survey will be published in The Sunflower Thank you. Please return by November 25, 1990 Atthur Lamey	THE PROPERTY OF THE PROPERTY O			Other (spec:	ty)	acres	
Dithane Apron (seed treatment) None used (list acres) Other (specify) Results of survey will be published in The Sunflower Thank you. Please return by November 25, 1990 Atthur Lamey							
Apron (seed treatment) None used (list acres) Other (specify) Results of survey will be published in The Sunflower Thank you. Please return by November 25, 1990 Author Lamey	3		3 200				
Results of survey will be published in The Sunflower Thank you. Please return by November 25, 1990 A Muy Arthur Lamey	w = 43		ns a rea no				
Results of survey will be published in The Sunflower Thank you. Please return by November 25, 1990 Author Lamey							
Results of survey will be published in The Sunflower Thank you. Please return by November 25, 1990 Atthur Lamey							
Thank you. Please return by November 25, 1990 Atthur Lamey		orner (specity)					
Thank you. Please return by November 25, 1990 Atthur Lamey		Decision of the control of the contr					
Thank you. Please return by November 25, 1990 Atthur Lamey	Desilts of survey will be an	blished in the constant					
Athur Lamey	Thank you. Please return his	November 25 1000					
	Tour Tour Targetti DA	1330			and the second		
	: 11						
	15/1	53					
	A NAUS CX MA	1011					
	· · · · · · · · · · · · · · · · · · ·					12	
xcension Plant Pathologist / /							
				2 0			

Figure 1. The survey form.

between oilseed and confection hybrids with regard to problems experienced, except that fewer disease problems were reported for oilseed than for confection hybrids. Worst production problems varied somewhat among the districts. The West Central and North West had more insect problems than other areas. By contrast, disease was more common in the North East and East Central districts (Table 3).

Table 1. Ranking of counties according to acres planted by respondents in 1990.

County	% of Respondents' Acresa	County	% of Respondents' Acres
Stutsman	11.7 ^b	McLean	2.0
Barnes	8.8 ^b	Eddy	1.9
Wells	6.2 ^b	Ward	1.9
LaMoure	6.1 ^b	Steele	1.7
Cass	5.6 ^b	McHenry	1.7
Foster	4.8 ^b	Bottineau	1.5
Ransom	4.8 ^b	Dickey	1.5
Benson	4.5 ^b	Grant	1.5
Ramsey	3.9	Morton	1.4
Pierce	3.0	Renville	1.3
Nelson	2.8	Towner	1.3
Cavalier	2.8	Griggs	1.2
Grand Forks	s 2.2	All others, ar	nd
Sargent	2.1	no response	11.8

aTotal acres planted by respondents=189,204

Insect Problems and Insecticide Use. Seed weevil was reported to be the No. 1 insect problem by 52% of respondents (Table 4). A higher percentage of confection growers, 66%, considered seed weevil to be the worst insect problem, as compared to 49% of oilseed growers. When the frequencies with which seed weevil was ranked No. 1, No. 2, and No. 3 were combined, seed weevil was reported to be one of the three worst insect problems by 76% of all respondents, 86% of confection growers and 74% of oilseed growers. Grasshoppers were cited as the worst insect problem by 25% of all respondents, 26% of oilseed growers and 19% of confection growers. Grasshoppers were one of the three worst insect problems for 61% of all respondents. Stem weevil was the third most commonly cited problem; it was the worst insect problem for 10% of all

Table 2. Worst sunflower production problem in 1990.

	Oil		All
Problem	Seed	Confection	Sunflower
	9	6 of responde	nts
Insects	22.1	22.9	22.6
Diseases	18.6	21.5	19.3
Weather	18.1	15.2	16.6
Bird Damage	16.9	15.7	16.3
Weeds	12.4	10.8	12.0
Emergence/Stand	4.1	6.7	4.9
Other	1.2	2.2	1.8
None	6.6	4.9	6.5

Table 3. Worst production problem in 1990, as ranked by respondents in each district.^a

Problem	State- wide	North West	North Central	North East	West Central	Central	East Central	South ^b West	South ^b Central	South East
				%	of respor	ndents				
Insects	22.6	29.7	26.4	12.1	54.2	27.1	16.9	60.0	13.3	21.8
Diseases	19.3	8.1	15.3	25.0	-	18.6	24.2	-	6.7	21.8
Weather	16.6	24.3	20.8	22.6	29.2	9.3	11.3	20.0	53.3	11.5
Bird Damage	16.3	-	16.7	11.3	4.2	26.4	15.3	-	1 <mark>3.3</mark>	20.7
Weeds	12.0	29.7	15.3	15.3	4.2	7.8	10.5	20.0	13.3	6.9
Emergence/Stand	4.9	-	4.2	2.4	8.3	4.7	6.5	=	-	10.3
Other	1.8	5.4	-	4.8	1 200	1.6	0.8	-	-	-
None	6.5	2.7	1.4	6.5	97	4.7	14.5	-	-	6.9

^aDistricts as defined by the North Dakota Agricultural Statistical Service.

^bTop 8 counties account for 52.5% of acres planted by respondents.

bLess than 20 respondents

respondents and one of the three worst insect problems for 39% of all respondents. Stem weevil was more frequently cited as the worst insect problem by oilseed growers (11%) than by confection growers (3%). No major insect problem was reported by 8% of respondents. Any other insect listed was of major importance to less than 2% of respondents.

Respondents in all districts ranked the seed weevil as the worst insect problem (Table 5). However, grasshoppers were considered to be nearly as great a problem as the seed weevil in the North East. Stem

weevils were considered to be fairly important in the West Central, South Central and South East districts. The seed weevil was also ranked as one of the three worst insect problems by growers in all districts but the North East, where grasshoppers were listed as one of the three worst insect problems (Table 6). Grasshoppers were almost as frequently listed as one of the worst three insect problems as seed weevils in the North Central district. Stem weevils were also considered to be very important insect pests in the West Central, South West, and South Central districts.

Table 4. Worst insect problem in 1990

	Sur	All nflower	Oil	Seed	Confection	
Insect	No. 1 Problem	No. 1,2, or 3 Problem	No. 1 Problem	No. 1,2, or 3 Problem	No. 1 Problem	No. 1,2, or 3 Problem
W W			% of re	spondents		
Seed weevil	51.8	75.7	48.8	74.1	65.6	85.5
Grasshopper	25.0	61.3	26.3	61.8	18.9	57.3
Stem weevil	9.6	39.1	11.3	43.2	3.1	
Sunflower beetle	1.9	13.2	1.5	11.1	3.1	32.2
Sunflower head moth	1.6	5.5	1.2	5.4	2.2	17.6
Cutworm	1.4	3.6	1.7	3.6	1.3	6.6
Sunflower midge	0.6	3.1	0.8	3.3	3000	4.4
Banded sunflower moth	0.3	1.6	0.0	1.2	0.4	3.1
None	7.7	7.7	8.3	8.3	0.4 4.8	2.6 4.8

Table 5. Worst insect problem in 1990, as ranked by respondents in each district.^a

				A PARTY OF THE PAR		Caci ais	tilot.			
Insect	State- Wide	North West	N <mark>orth</mark> Central	North East	West Central	Central	East Central ^b	South West ^b	South Central	South East
					% of resp	ondents				
Seed weevil	51.8	60.5	43.1	38.0	58.3	51.5	53.2	80.0	66.7	70.1
Grasshopper	25.0	23.7	36.1	37.2	20.8	26.2	22.6	20.0	6.7	3.4
Stem weevil	9.6		13.9	3.9	20.8	9.2	8.1		20.0	17.2
Sunflower beetle	1.9	7	1.4	6.2		0.8	0.8		20.0	11.2
Sunflower head moth	1.6	2.6	2.8	1.6		1.5	1.6		_	
Cutworm	1.4	2.6	1.4	14		2.3	0.8	1000	· • <u>· · · · · · · · · · · · · · · · · ·</u>	3.4
Sunflower midge	0.6	p 11 2	1.4	1.6	-	44	0.8			5.4
Banded sunflower moth	0.3	. 3.		-	The second second	0.8	0.8			19 200
None	7.7	10.5	2 2 1	11.6		7.7	11.3	2 <u>2</u>	6.7	5.7

^aDistricts as defined by the North Dakota Agricultural Statistical Service.

bLess than 20 respondents

Table 6. Insects ranked as one of three worst insect problems in 1990, according to respondents in each district.

			400				- 40	1.0		
Insect	State- Wide	North West	North Central	North . East	West Central	Central	East Central	South West ^b	South Central ^b	South East
4 4 5 4 5 2	41 12				- % of re	sponden	ts		Ğ	
Seed weevil	75.7	78.9	76.4	61.2	83.3	78.5	72.6	100.0	93.3	88.5
Grasshopper	61.3	52.6	75.0	65.1	83.3	67.7	71.8	40.0	66.7	37.9
Stem weevil	39.1	44.7	40.3	14.0	70.8	45.4	36.3	60.0	66.7	55.2
Sunflower beetle	13.2	13.2	18.1	26.4	4.2	12.3	8.1		a #	3.4
Sunflower head moth	5.5	5.3	8.3	6.2	or a	6.2	5.6		* -	2.3
Cutworm	3.6	2.6	6.9	-	× 5%	6.2	1.6		6.7	6.9
Sunflower midge	3.1	2.6	4.2	3.1	-	-	4.8	-	-	4.6
Banded sunflower moth	1.6	-	-	2.3	_	2.3	1.6	9 1	=	1.1
None	7.7	10.5	=:	11.6	48	7.7	11.3	89 2 8	6.7	5.7

^aDistricts as defined by the North Dakota Agricultural Statistical Service.

Parathion was the most commonly used insecticide. When data for all types of parathion (methyl, 6-3, and ethyl) were combined, parathion was sprayed on 64% of respondents' acres (Table 7). A greater percentage of confection acres than oilseed acres was sprayed with insecticide. Asana XL was the second most commonly used insecticide, applied on 15% of respondents' acres, followed by Furadan (15G and 4F), applied on 7% of respondents' acres. Less than 2% of respondents' acres were treated with any other insecticide.

Respondents did not treat 29% of their oilseed acres or 9% of their confection acres.

Disease Problems and Fungicide Use. Sclerotinia wilt was the worst disease problem for 35% of respondents and it was one of the three worst disease problems for 49% of respondents (Table 8). No disease problem was reported by 32% of respondents. Phoma black stem was cited as the worst disease problem by 13% of respondents and as one of the three worst disease problems by 27% of respondents. Sclerotinia head rot, downy mildew and rust were also mentioned frequently as major disease problems; they were cited as one of the three worst disease problems by 27%. 17% and 21% of respondents respectively. Most disease problems were equally prevalent on both oilseed and confection hybrids; however, Phoma black stem was cited as the worst disease problem affecting oilseed hybrids by 15% of respondents and for confection hybrids by only 10% of respondents. Rust was more frequently cited as the worst or one of the three worst

Table 7. Insecticide use on sunflower in 1990.

Insecticide	All Sunflowe <mark>r</mark>	Oil Seed	Confection
in .	% of r	esponden	ts' acres
Methyl parathion	34.9	35.0	70.3
6-3 parathion	16.9	15.1	41.1
Asana XL	14.1	18.8	23.8
Ethyl parathion	12.3	13.7	26.7
Furadan 15G	- 4.1	4.6	5.9
Furadan 4F	2.4	3.0	4.1
Lindane/maneb	1.4	2.0	0.9
Others	0.9	1.2	1.0
Sevin XLR Plus	0.9	1.3	0.7
Lorban 4E	0.3	0.2	0.8
None used	21.1	29.4	9.3
Total parathion	64.1	63.8	138.1
Total Furadan	6.5	7.6	10.0

disease problems affecting confection hybrids (12% and 37% of respondents) than for oilseed hybrids (3% and 17% of respondents).

Sclerotinia wilt was most frequently ranked as the worst disease problem by respondents in most cropping districts, but it was more frequently cited as a problem in the North East, Central, and East Central

bLess than 20 respondents

districts (Table 9). Sclerotinia wilt was also cited as one of the three worst disease problems by respondents in many cropping districts, but Phoma black stem and Sclerotinia head rot also were frequently cited (Table 10). Phoma black stem was cited as one of the three worst disease problems by over 30% of respondents in the North Central and Central districts.

The most frequently used fungicide was Apron seed treatment, used on 49% of all respondents' acres (Table 11). Dithane fungicide was used for rust control on 1% of all of respondents' acres; however it was

used on only 0.3% of respondents' oilseed acres and on 3.3% of respondents' confection acres.

Bird Damage. Approximately two-thirds of all respondents estimated that they had 0-5% loss from birds; the other one-third had more than 5% as follows: 19% claimed losses of 5-10% and 14% estimated losses to be greater than 10% (Table 12). Bird losses appeared to be higher than the statewide average in the North Central and Central cropping districts, where over 22% of respondents reported losses in excess of 10% (Table 13). Bird losses were approximately the same in oilseed and confection hybrids.

Table 8. Worst disease problem in 1990.

		All flower	Oil	Seed	Cor	fection
Disease	No. 1 Problem	No. 1,2, or 3 Problem	No. 1 Problem	No. 1,2, or 3 Problem	No. 1 Problem	No. 1,2, or 3 Problem
		9	6 of respon	ndents		
Sclerotinia wilt	34.8	49.3	32.5	47.1	43.3	58.7
Phoma black stem	13.1	26.5	15.4	28.1	9.6	30.3
Sclerotinia head rot	7.5	26.6	8.1	27.3	6.3	27.9
Downy mildew	6.3	16.6	6.3	16.0	5.3	15.4
Rust	5.8	20.6	3.3	16.7	12.0	37.0
Other :	0.9	1.5	1.0	1.9	0.5	1.0
None	31.6	31.6	33.3	33.3	23.1	23.1
Sclerotinia TOTAL	42.3	75.9	40.6	74.4	49.6	86.6

Table 9. Worst disease problem in 1990, as ranked by repondents in each district.^a

Disease	State- Wide	North West	North Central	North East	West Central	Central	East Central	South West ^b	South Central ^b	South East
7 V.					% of res	pondent	s			
Sclerotinia wilt	34.8	16.1	28.4	40.8	10.0	41.0	40.5	20.0	1240	32.5
Phoma black stem	13.1	9.7	20.9	4.2	10.0	19.7	5.0	_	15.4	23.4
Sclerotinia head rot	7.5	6.5	6.0	10.0	. <u>.</u>	6.6	8.3	ne the	15.4	6.5
Downy mildew	6.3	_	6.0	9.2	€. 54	6.6	7.4	2 7. 5	7.7	5.2
Rust	5.8	3.2	7.5	10.8	5.0	3.3	5.8	w	-	3.9
Other	0.9		5E	ie status	5.0	0.8	8.0		10 M	2.6
None	31.6	64.5	31.3	25.0	70.0	22.1	32.2	80.0	61.5	26.0
Sclerotina TOTAL	42.3	22.6	34.4	50.8	10.0	47.6	48.8	20.0	15.4	39.0

^aDistricts as defined by the North Dakota Agricultural Statistics Service.

bLess than 20 respondents

Table 10. Diseases ranked as one of three worst disease problems in 1990, according to respondents in each district.2

State- Wide	North West	North Central	North East	West Central	Central	East Central	South West ^b	South Central ^b	South East
				% of re	spondent	8			
49.3	16.1	47.8	55.0	10.0	60.7	50.4	20.0	23.1	50.6
26.5	25.8	31.3	15.8	20.0	35.2	18.2	20.0	15.4	40.3
26.6	22.6	16.4	25.0	10.0	30.3	30.6	-	30.8	30.0
16.6	6.5	20.9	20.0	5.0	17.6	16.5	0 .0	15.4	15.6
20.6	9.7	19.4	37.5	10.0	15.6	18.2		*	16.9
1.5	2 	220	, 0.8	5.0	2.5	1.7	1.75	-	2.6
31.6	64.5	31.3	25.0	70.0	22.1	32.2	80.0	61.5	26.0
75.9	38.7	64.2	80.0	20.0	91.0	81.0	20.0	53.9	80.6
	49.3 26.5 26.6 16.6 20.6 1.5 31.6	Wide West 49.3 16.1 26.5 25.8 26.6 22.6 16.6 6.5 20.6 9.7 1.5 - 31.6 64.5	Wide West Central 49.3 16.1 47.8 26.5 25.8 31.3 26.6 22.6 16.4 16.6 6.5 20.9 20.6 9.7 19.4 1.5 - - 31.6 64.5 31.3	Wide West Central East 49.3 16.1 47.8 55.0 26.5 25.8 31.3 15.8 26.6 22.6 16.4 25.0 16.6 6.5 20.9 20.0 20.6 9.7 19.4 37.5 1.5 - - , 0.8 31.6 64.5 31.3 25.0	Wide West Central East Central 49.3 16.1 47.8 55.0 10.0 26.5 25.8 31.3 15.8 20.0 26.6 22.6 16.4 25.0 10.0 16.6 6.5 20.9 20.0 5.0 20.6 9.7 19.4 37.5 10.0 1.5 - , 0.8 5.0 31.6 64.5 31.3 25.0 70.0	Wide West Central East Central Central 49.3 16.1 47.8 55.0 10.0 60.7 26.5 25.8 31.3 15.8 20.0 35.2 26.6 22.6 16.4 25.0 10.0 30.3 16.6 6.5 20.9 20.0 5.0 17.6 20.6 9.7 19.4 37.5 10.0 15.6 1.5 - - 0.8 5.0 2.5 31.6 64.5 31.3 25.0 70.0 22.1	Wide West Central East Central Central Central 49.3 16.1 47.8 55.0 10.0 60.7 50.4 26.5 25.8 31.3 15.8 20.0 35.2 18.2 26.6 22.6 16.4 25.0 10.0 30.3 30.6 16.6 6.5 20.9 20.0 5.0 17.6 16.5 20.6 9.7 19.4 37.5 10.0 15.6 18.2 1.5 - - 0.8 5.0 2.5 1.7 31.6 64.5 31.3 25.0 70.0 22.1 32.2	Wide West Central East Central Central Central Westb 49.3 16.1 47.8 55.0 10.0 60.7 50.4 20.0 26.5 25.8 31.3 15.8 20.0 35.2 18.2 20.0 26.6 22.6 16.4 25.0 10.0 30.3 30.6 - 16.6 6.5 20.9 20.0 5.0 17.6 16.5 - 20.6 9.7 19.4 37.5 10.0 15.6 18.2 - 1.5 - - 0.8 5.0 2.5 1.7 - 31.6 64.5 31.3 25.0 70.0 22.1 32.2 80.0	Wide West Central East Central Central Central Westb Centralb 49.3 16.1 47.8 55.0 10.0 60.7 50.4 20.0 23.1 26.5 25.8 31.3 15.8 20.0 35.2 18.2 20.0 15.4 26.6 22.6 16.4 25.0 10.0 30.3 30.6 - 30.8 16.6 6.5 20.9 20.0 5.0 17.6 16.5 - 15.4 20.6 9.7 19.4 37.5 10.0 15.6 18.2 - - 1.5 - - 0.8 5.0 2.5 1.7 - - 31.6 64.5 31.3 25.0 70.0 22.1 32.2 80.0 61.5

^aDistricts as defined by the North Dakota Agricultural Statistics Service

Table 11. Fungicide use on sunflower in 1990.

WAR PERSON		Table 12. Of	op 1033 dae to b	iru damage	11 1330.
Oil			All	Oil	
Seed	Confection	% Loss	Sunflower	Seed	Confection

65.2 20.1 10.3 4.5

Fungicide	All Sunflower	Oil Seed	Confection	% Loss	All Sunflower	Oil Seed	(
Ä	% of re	sponden	its' acres		%	of responder	nts -
Apron (seed treatment)	48.5	60.5	64.8	0-5	66.3	65.4	
Dithane	1.3	0.3	3.3	5-10	19.3	19.7	
Others	1.0	1.1	1.6	10-25	10.2	10.5	
None	32.1	41.2	38.2	25- 5 0	3.5	3.5	
				50-100	0.2	0.8	

Table 13. Crop loss due to bird damage in 1990, according to respondents in each district.^a

% Loss	State- Wide	North West	North Central	North East	West Central	Central	East Central	South West ^b	South Central ^b	South East
					%(of respond	ents			
0-5	66.3	73.7	50.7	79.2	73.9	53.5	75.4	40.0	61.5	60.9
5-10	19.3	13.2	26.8	16.0	26.1	22.8	15.1	60.0	23.1	19.5
10-25	10.2	7.9	16.9	4.0	7.	15.0	7.1	*	15.4	13.8
25-50	3.5	5.3	2.8	0.8		7.9	2.4			4.6
50-100	0.2	-	2.8			0.8		2	ne sto	1.1

^aDistricts as defined by the North Dakota Agricultural Statistics Service.

bLess than 20 repondents

bLess than 20 respondents

Weed Problems and Herbicide Use. The four worst weed problems listed in order of severity were kochia, foxtail (pigeongrass), Russian thistle and wild mustard. Kochia was cited as the worst weed problem by 42% of respondents and as one of the three worst weed problems by 66% of respondents (Table 14). Foxtail was cited as the worst weed problem by 17% of respondents and one of the three worst weed problems by 41% of respondents. Russian thistle was cited as the worst weed problem by 9% of respondents and as one of the three worst weed problems by 30% of respon-

dents. Wild mustard was cited as the worst weed problem by 6% of respondents and as one of the three worst weed problems by 21% of respondents. Most other weeds were cited by only a small percentage of respondents. Growers of both oilseed and confection hybrids ranked the weeds in the same order of importance.

Kochia was more frequently cited as the worst weed problem by respondents in the North Central and West Central districts, where it was cited by over 50% of respondents (Table 15). Russian thistle was cited as

Table 14. Worst weed problem in 1990.

52		All flower		Oil eed	Confection	
Weed	No. 1 Problem	No. 1,2, or 3 Problem	No. 1 Problem	No.1,2, or 3 Problem	No. 1 Problem	No.1,2, or 3 Problem
54			****			
Kochia	42.3	66.2	43.1	66.9	40.4	66.2
Foxtail (pigeongrass)	16.7	41.2	16.9	42.7	16.2	38.6
Russian thistle	9.1	30.1	9.8	30.5	6.1	28.5
Wild mustard	6.2	20.8	5.6	20.3	9.6	23.7
Others	2.5	4.5	2.4	4.5	2.6	5.3
Cocklebur	2.3	4.2	1.7	3.0	3.5	6.6
Canada thistle	2.3	3.4	2.1	3.2	4.4	6.1
Redroot pigweed	1.6	8.2	1.5	8.3	1.3	7.5
Wild oats	1.6	11.3	1.5	11.1	1.3	12.3
Wild buckwheat	8.0	3.3	0.6	3.4	0.9	3.1
None	14.6	14.6	14.7	14.7	13.6	13.6

Table 15. Worst weed problem in 1990, as ranked by respondents by district.^a

Weed	State- Wide	North West	North Central	North East	West Central	Central	East Central	South West ^b	South Central ^b	South East
-					AND THE PROPERTY OF THE PROPER	spondents	APCDOONSE HERVING VICE		Contrai	
Kochia Foxtail	42.3	42.5	58.1	38.3	50.0	39.1	36.5	80.0	60.0	38.2
(pigeongrass)	16.7	15.0	16.2	8.6	25.0	16.5	22.2	_	6.7	21.3
Russian thistle	9.1	22.5	6.8	8.6	8.3	12.8	3.2	.50	13.3	10.1
Wild mustard	6.2	7.5	6.8	12.5		2.3	7.9	() ()	(100 Mileseles)	2.2
Others	2.5	2.5	-	2.3	, 77 8	3.8	3.2	0	6.7	2.2
Cocklebur	2.3	-	2.7	2.3	-	5.3	1.6	 c		1.1
Canada thistle	2.3	5.0	1.4	4.7	-	1.5	0.8	<u> 1860 lin</u>	<u>-</u>	3.4
Redroot pigweed	1.6	-	1425 M	0.8	4.2	0.8	3.2	-	P. 60	3.4
Wild oats	1.6	2.5	2.7	3.1	1245 1245	2.3	4	Lady	<u>=</u>	_
Wild buckwheat	0.8	4	1.4	-	4.2	1.5	0.8		1044 1044	
None	14.6	2.5	4.1	18.8	8.3	14.3	20.6	20.0	13.3	18.0

^aDistricts as defined by the North Dakota Agricultural Statistics Service

bLess than 20 respondents

the worst weed problem by 23% of respondents in the North West district, well above the statewide average of 9%. Wild mustard was a more common weed problem in the North East, where it was cited by 13% of respondents, compared to 6% of respondents in the rest of the state. Kochia was cited as one of the three worst weed problems by over 80% of respondents in the North Central, South West and South Central districts (Table 16). Russian thistle was cited as one of the three worst weed problems by over 50% of the respondents in the North West and South Central districts.

The two dinitro analine herbicides Treflan and Sonalan accounted for the majority of herbicide use on sunflower (Table 17). The combined spring and fall application of Treflan plus application of Sonalan accounted for 96% of respondents' acres. Spring-applied Treflan accounted for 51% of respondents' acres, Sonalan for 36%, and fall-applied Treflan for 9%. Weed control methods other than chemicals used by respondents included cultivation and hand weeding. Cultivation was used by 89% of respondents. Most respondents used one cultivation, but a few used more than one (see discussion under alternative control measures).

Table 17. Use of herbicides and alternative control methods in 1990.

- The Eventue of the Committee of the Co	2 2					
9	All Sur	nflower				
Herbicide	% of respondents	% of respondents' acres				
Treflan (spring)	58.7	51.2				
Sonalan	43.3	36.2				
Treflan (fall)	11.0	9.0				
Roundup (preplant)	4.6	1.2				
Assert	3.2	1.1				
Poast	2.9	0.6				
Prowl (spring)	1.7	0.8				
Lasso	0.6	0.4				
Treflan + Eptam/Genep	0.5	0.2				
Sonalan + Eptam/Genep	0.3	0.3				
Leafex-3/Defol	0.3	0.1				
TOTAL:	127.1	101.1				
	All Su	ınflower				
Other Weed Control	% of respondents	The True section for the section of				
		acres				

	All Sunflower				
Other Weed Control	% of respondents	% of respondents' acres			
Cultivation	88.8	115.4			
Hand weeding	0.6	0.5			
Other	1.4	0.9			

Table 16. Weeds ranked as one of three worst weed problems in 1990, according to respondents in each district.^a

Weed	State- Wide	North West	North Central	North East	West Central	Central	East Central	South West ^b	South Central ^b	South East
~ T					% of respo	ndents				
Kochia	66.2	67.5	81.1	63.3	66.7	65.4	59.5	80.0	80.0	62.9
Foxtail										
(pigeongrass)	41.2	37.5	40.5	25.0	66.7	42.1	41.3	60.0	46.7	51.7
Russian thistle	30.1	52.5	35.1	28.1	29.2	31.6	15.1	20.0	53.3	36.0
Wild mustard	20.8	15.0	25.7	28.1	8.3	21.1	22.2	X = 1	20.0	12.4
Others	4.5	12.5	1.4	3.9	12	6.8	3.2	-	6.7	4.5
Cocklebur	4.2	· ·	2.7	5.5	-	8.2	4.0	5. -	1984	2.2
Canada thistle	3.4	5.0	4.1	7.0	/ 	1.5	2.4	(=	5 %	4.5
Redroot pigweed	8.2	5.0	5.4	6.3	8.3	6.8	14.3	-	6.7	9.0
Wild oats	11.3	20.0	6.8	14.8	8.3	9.8	15.9	20.0	-	2.2
Wild buckwheat	3.3	5.0	9.5	1.6	4.2	3.0	0.8	(-	-	4.5
None	14.6	2.5	4.1	18.8	8.3	14.3	20.6	20.0	13.3	18.0

^aDistricts as defined by the North Dakota Agricultural Statistical Service.

bLess than 20 repondents

Respondents ranked most herbicides as providing good to excellent weed control (Table 18). Herbicides highly ranked included Treflan (spring and fall), Sonalan, Roundup (preplant), Assert, and Poast. Prowl (spring) and Lasso were less frequently ranked as giving good to excellent weed control. All herbicides except Prowl caused none to slight injury on sunflower; Prowl was cited by 22% of respondents as causing moderate sunflower injury. Cultivation was ranked as giving good to excellent weed control by 81% of respondents, but 25% of respondents cited cultivation as causing moderate sunflower injury.

Alternative Control Measures. Crop Consultants. Only 11% of respondents reported that they hired a crop consultant in 1990 (Table 19). Of the respondents who hired a crop consultant, 19% reported no economic gain from the use of a crop consultant, 33% reported an economic gain of less than \$10/acre, and 48% reported an economic return of \$10/acre or more: 38% reported an economic gain of \$10-20/acre and 10% reported an economic gain of over \$20/acre (Table 20).

Table 19. Use of a hired crop consultant in 1990.

588		% of	7
	Consultant	Respondents	- 1 - 4 s
10 0	YES	11.4	
	NO	88.6	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1

Table 20. Economic return from using a crop consultant in 1990.

	Economic Return		% of Respondents		7
	\$0/A	1	19.0	Ø.	, Att 6
	\$0-5/A		17.5	Ġ.	
	\$5-10/A		15.9		
	\$10-15/A		25.4		2 5
	\$15-20/A		12.7	$p_{i_1}^{(2)} = n$	0.00
	\$20 +/A		9.5		
-					

Table 18. Effect of herbicides and alternative methods on weed control and injury in 1990.

		Weed C	ontrol			Sunflow	er Injury	24.
Herbicide	Excel.	Good	Fair	Poor	None	Slight	Moderate	Severe
				% of r	espondents	}		
Treflan (spring)	30	. 50	17	3	89	9	1	1
Sonalan	48	38	11	3	91	8		<u>.</u>
Treflan (fall)	25	48	18	9	93	7		* * *
Roundup (preplant)	48	41	10		100	_	_	-
Assert	63	32	5		56	44		_
Poast	63	26	11		93	7		
Prowl (spring)	18	46	27	9	9	78	22	
Lasso	25	25	25	25	75	25		
Treflan +						E E		
Eptam/Genep	67		33		100	_		4
Sonalan +								7 2
Eptam/Genep	- 	100			100		- 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1	
Leafex-3/Defol		100			100			-
Other weed				i a jar		4		The state of
Control Methods						***		
Cultivation	31	50	18	2	42	54	25	
Hand weeding	50	25	25		25	50	25	
No herbicide		50		50	50		50	Maria de la companya
Other	17	33	17	33	20	20		60

Crop Rotation was listed as an alternative pest control measure by 66% of respondents. Wheat was grown as the preceding crop by 71% of respondents, barley by 21% and oats by 3% (Table 21). Respondents commonly had grown sunflower three, four or five years prior to the 1990 sunflower crop (Table 22). Respondents indicated that four years elapsed since the previous sunflower crop in 36% of responses, three years in 29% of responses, and five years in 15% of responses. Only 6% of respondents indicated that sunflower had been grown 0, one, or two years prior to the 1990 sunflower crop.

Cultivation was used by most respondents. A single cultivation was used by 69% of respondents, two cultivations by 20%, three cultivations by 5% and four cultivations by 1% (Table 23).

Resistant Varieties were used by 29% of respondents on 27% of acres in 1990. Only 1% of respondents reported "other alternate control practices", which were used on <1% of acres in 1990.

Table 21. Crop rotation: previous crop grown in 1990.a

	Preceding Crop	% of Respondents	
	Wheat	71.1	
	Barley	21.3	
	Oats	2.5	
	Corn	1.4	
¥3	Flax	0.8	
	Summer fallow	0.6	
	Rye	0.6	
	Sunflower	0.6	
	Alfalfa	0.4	
	Soybeans	0.4	
	Navy beans	0.2	
	Buckwheat	0.2	

^a66.0% of respondents listed crop rotation as an alternative pest control measure; crop rotation was listed for 67.9% of respondents' acres.

Table 22. Crop rotation: in 1990: years since previous sunflower crop.

	Years	% of Respondents	
	0	0.2	
	market and the	0.3	
	2	5.9	to jake a
/	3	28.5	
	4	36.3	
The state of	5	14.5	
	6	4.3	
	7	1.9	
	8	2.1	
	9	0.9	
	10	1.6	
	13	0.2	
	No previous		
	sunflower crop	3.3	

Table 23. Number of cultivations used in 1990.

	No. of Cultivat <mark>i</mark> ons	% of Respondents	
	0	4.4	
	1	68.8	
	2	20.1	
S	3	5.4	
	4	1.3	

LITERATURE CITED

1. North Dakota Agricultural Statistics Service. 1991. North Dakota Agricultural Statistics. 1991. 101 p.

Helping You Put Knowledge To Work